




---

## Product Datasheet

### SRD5A1 Antibody (orb1246911)

<b>Description</b>	SRD5A1 Antibody	
<b>Species/Host</b>	Goat	
<b>Reactivity</b>	Human	
<b>Conjugation</b>	Unconjugated	
<b>Tested Applications</b>	ELISA, IHC	orb1246911 (4 ug/ml) staining of paraffi...
<b>Immunogen</b>	The immunogen for this antibody is: ATATGVAEERLLC	
<b>Target</b>	SRD5A1	
<b>Preservatives</b>	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.	
<b>Form/Appearance</b>	Liquid	
<b>Concentration</b>	500 ug/mL	
<b>Storage</b>	Aliquot and store at -20°C. Minimize freezing and thawing.	
<b>Note</b>	For research use only	
<b>Application notes</b>	Peptide ELISA: antibody detection limit dilution 1:128000. Immunohistochemistry: In paraffin embedded Human Prostate shows reticulate staining in the cytoplasm of secretory cells in the gland. Recommended concentration: 4-6ug/ml. Paraffin embedded Human Liver. Recommended concentration: 5ug/ml.	 orb1246911 (5 ug/ml) staining of paraffi...
<b>Clonality</b>	Polyclonal	
<b>Uniprot ID</b>	<a href="#">P18405</a>	
<b>NCBI</b>	<a href="#">NP_001038.1</a>	
<b>Dilution Range</b>	Peptide ELISA: antibody detection limit dilution 1:128000. Immunohistochemistry: In paraffin embedded Human Prostate shows reticulate staining in the cytoplasm of secretory cells in the gland. Recommended concentration: 4-6ug/ml. Paraffin embedded Human Liver. Recommended concentration: 5ug/ml.	
<b>Expiration Date</b>	12 months from date of receipt.	