



Product Datasheet

NRTN Antibody (orb1247205)

| | | |
|----------------------------|--|---|
| Description | NRTN Antibody |  |
| Species/Host | Goat |  |
| Reactivity | Human | orb1247205 (0.1 ug/ml) staining of Human... |
| Conjugation | Unconjugated | |
| Tested Applications | ELISA, IHC, WB | |
| Immunogen | The immunogen for this antibody is: C-DAHSRYHTVHELASRE | |
| Target | NRTN | |
| Preservatives | Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing. |  |
| Form/Appearance | Liquid | orb1247205 (5 ug/ml) staining of paraffi... |
| Concentration | 500 ug/mL | |
| Storage | Aliquot and store at -20°C. Minimize freezing and thawing. | |
| Note | For research use only | |
| Application notes | Peptide ELISA: antibody detection limit dilution 1:64000. Western Blot: Approx 25kDa band observed in Human Heart lysates (calculated MW of 22.4kDa according to NP_004549.1). Recommended concentration: 0.1-0.3ug/ml. Primary incubation was 1 hour. Immunohistochemistry: Paraffin embedded Human Colon (Submucosal Plexus). Recommended concentration: 5ug/ml. | |
| Clonality | Polyclonal | |
| Uniprot ID | Q99748 | |
| NCBI | NP_004549.1 | |
| Dilution Range | Peptide ELISA: antibody detection limit dilution 1:64000. Western Blot: Approx 25kDa band observed in Human Heart lysates (calculated MW of 22.4kDa according to NP_004549.1). Recommended concentration: 0.1-0.3ug/ml. Primary incubation was 1 hour. Immunohistochemistry: Paraffin embedded Human Colon (Submucosal Plexus). Recommended concentration: 5ug/ml. | |
| Expiration Date | 12 months from date of receipt. | |