
Product Datasheet

NPY5R Antibody (orb1247207)

Description

NPY5R Antibody

Species/Host

Goat

Reactivity

Human

Conjugation

Unconjugated

Tested

ELISA, IHC, IP, WB

Applications
Immunogen

The immunogen for this antibody is: C-GLSNKENRLEEN

Target

NPY5R

Preservatives

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

Form/Appearance

Liquid

Concentration

500 ug/mL

Storage

Aliquot and store at -20°C. Minimize freezing and thawing.

Note

For research use only

Application notes

Peptide ELISA: antibody detection limit dilution 1:32000. Western Blot: Approx 50kDa band observed in Human Brain, Breast and Duodenum lysates (calculated MW of 50.7kDa according to NP_006165.1). Recommended concentration: 0.3-1ug/ml. Primary incubation was 1 hour. Immunohistochemistry: In paraffin embedded Human Intestine shows staining of ganglion cells. Recommended concentration, 4-6ug/ml. Immunoprecipitation: Customer feedback showed specific results when this product was used in IP.

Clonality



Polyclonal

Uniprot ID
[Q15761](#)
NCBI
[NP_006165.1](#)
Dilution Range

Peptide ELISA: antibody detection limit dilution 1:32000. Western Blot: Approx 50kDa band observed in Human Brain, Breast and Duodenum lysates (calculated MW of 50.7kDa according to NP_006165.1). Recommended concentration: 0.3-1ug/ml. Primary incubation was 1 hour. Immunohistochemistry: In paraffin embedded Human Intestine shows staining of ganglion cells. Recommended concentration, 4-6ug/ml. Immunoprecipitation: Customer feedback showed specific results when this product was used in IP.

Expiration Date

12 months from date of receipt.


 orb1247207
(0.3 ug/ml)
staining of
Human...

 orb1247207
(3.8 ug/ml)
staining of
paraf...