

Product Datasheet

EXOC7 Antibody (orb1247540)

Description	EXOC7 Antibody	
Species/Host	Goat	
Reactivity	Human	
Conjugation	Unconjugated	
Tested Applications	ELISA, IHC, WB	orb1247540 (0.5 ug/ml) staining of Human...
Immunogen	The immunogen for this antibody is: C-DNIKNDPDKEYN	
Target	EXOC7	
Preservatives	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.	
Form/Appearance	Liquid	
Concentration	500 ug/mL	
Storage	Aliquot and store at -20°C. Minimize freezing and thawing.	orb1247540 (0.5 ug/ml) staining of paraf...
Note	For research use only	
Application notes	Peptide ELISA: antibody detection limit dilution 1:16000. Western Blot: Approx 70kDa band observed in Human Lung lysates (calculated MW of 74.7kDa according to NP_056034.2). Recommended concentration: 0.5-1.5ug/ml. Immunohistochemistry: In paraffin embedded Human Adrenal Gland shows staining of cytoplasm membranes and cytoplasm of secretory cells in the medulla . Recommended concentration, 5-10ug/ml.	
Clonality	Polyclonal	
Uniprot ID	Q9UPT5	
NCBI	NP_001013861.1 , NP_001138770.1 , NP_001138771.1 , NP_056034.2 , NP_001138769.1	
Dilution Range	Peptide ELISA: antibody detection limit dilution 1:16000. Western Blot: Approx 70kDa band observed in Human Lung lysates (calculated MW of 74.7kDa according to NP_056034.2). Recommended concentration: 0.5-1.5ug/ml. Immunohistochemistry: In paraffin embedded Human Adrenal Gland shows staining of cytoplasm membranes and cytoplasm of secretory cells in the medulla . Recommended concentration, 5-10ug/ml.	
Expiration Date	12 months from date of receipt.	