



---

## Product Datasheet

### DCXR Antibody (orb1247608)

<b>Description</b>	DCXR Antibody	
<b>Species/Host</b>	Goat	
<b>Reactivity</b>	Human	
<b>Conjugation</b>	Unconjugated	
<b>Tested Applications</b>	ELISA, IHC, WB	orb1247608 staining (0.1 ug/ml) of human...
<b>Immunogen</b>	The immunogen for this antibody is: GSTLPVEGGFWAC	
<b>Target</b>	DCXR	
<b>Preservatives</b>	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.	
<b>Form/Appearance</b>	Liquid	
<b>Concentration</b>	500 ug/mL	
<b>Storage</b>	Aliquot and store at -20°C. Minimize freezing and thawing.	orb1247608 (1 ug/ml) staining of paraffi...
<b>Note</b>	For research use only	
<b>Application notes</b>	Peptide ELISA: antibody detection limit dilution 1:32000. Western Blot: Approx 26kDa band observed in human kidney, liver and testes lysates (calculated MW of 25.9kDa according to NP_057370.1). Recommended for use at 0.1-0.3ug/ml. Immunohistochemistry: In paraffin embedded human kidney, stains membrane / cytoplasm in epithelial cells of renal glomeruli and tubules. Recommended concentration, 1-2ug/ml.	
<b>Clonality</b>	Polyclonal	
<b>Uniprot ID</b>	<a href="#">Q7Z4W1</a>	
<b>NCBI</b>	<a href="#">NP_057370.1</a>	
<b>Dilution Range</b>	Peptide ELISA: antibody detection limit dilution 1:32000. Western Blot: Approx 26kDa band observed in human kidney, liver and testes lysates (calculated MW of 25.9kDa according to NP_057370.1). Recommended for use at 0.1-0.3ug/ml. Immunohistochemistry: In paraffin embedded human kidney, stains membrane / cytoplasm in epithelial cells of renal glomeruli and tubules. Recommended concentration, 1-2ug/ml.	
<b>Expiration Date</b>	12 months from date of receipt.	