



Product Datasheet

CARD6 Antibody (orb1247705)



www.biorbyt.com

Descriptionnts. CARD6 Antibody

Species/Host Goat

Reactivity Human

Conjugation Unconjugated

Tested ELISA, IHC

Applications

Immunogen The immunogen for this antibody is: ATESTPSEIIERER-C

Target CARD6

Preservatives Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH

7.3 with 0.5% bovine serum albumin. Aliquot and store at -

20°C. Minimize freezing and thawing.

Form/Appearance Liquid

Concentration 500 ug/mL

Storage Aliquot and store at -20°C. Minimize freezing and thawing.

Note For research use only

Application notes Peptide ELISA: antibody detection limit dilution

1:32000.Western Blot:Approx. 45-50kDa and 25-30kDa bands observed in A549, Human Heart and Human Skeletal Muscle lysates at 1ug of primary. Please note that the bands we observe are different from the predicted size of approx. 119kDa according to NP_115976. Currently we cannot find an explanation for this in the literature. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any splice

variants/modified forms been

reported?Immunohistochemistry:In paraffin embedded Human Pancreas shows variable staining of cytoplasm in acinar cells. Recommended concentration, 5-10ug/ml.

Clonality Polyclonal

Uniprot ID Q9BX69

NCBI NP_115976.2

Dilution Range Peptide ELISA: antibody detection limit dilution

1:32000.Western Blot:Approx. 45-50kDa and 25-30kDa bands observed in A549, Human Heart and Human Skeletal Muscle lysates at 1ug of primary. Please note that the bands we observe are different from the predicted size of approx. 119kDa according to NP_115976. Currently we cannot find an explanation for this in the literature. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any splice

orb1247705 (5 ug/ml) staining of paraffi...