




Product Datasheet

SFN Antibody (orb1247868)

Description	SFN Antibody	
Species/Host	Goat	
Reactivity	Human, Mouse, Rat	
Conjugation	Unconjugated	
Tested Applications	ELISA, IHC, WB	orb1247868 (0.03 ug/ml) staining of Huma...
Immunogen	The immunogen for this antibody is: C-DLHTLSEDSYKDST	
Target	SFN	
Preservatives	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.	
Form/Appearance	Liquid	
Concentration	500 ug/mL	
Storage	Aliquot and store at -20°C. Minimize freezing and thawing.	orb1247868 (2 ug/ml) staining of paraffi...
Note	For research use only	
Application notes	Peptide ELISA: antibody detection limit dilution 1:16000. Western Blot: Approx 26-28kDa band observed in Human, Mouse and Rat Skin lysates and in lysates of cell lines A431 and U2OS (calculated MW of 27.8kDa according to Human NP_006133.1, Mouse NP_061224.2 and Rat XP_003750096.1). Recommended concentration: 0.03-0.1ug/ml. Primary incubation 1 hour at room temperature. Immunohistochemistry: In paraffin embedded Human Skin shows strong nuclear staining in selected cells of the basal layer. Recommended concentration, 2-4ug/ml.	
Clonality	Polyclonal	orb1247868 (0.03 ug/ml) staining of A431...
Uniprot ID	P31947	
NCBI	NP_006133.1	
Dilution Range	Peptide ELISA: antibody detection limit dilution 1:16000. Western Blot: Approx 26-28kDa band observed in Human, Mouse and Rat Skin lysates and in lysates of cell lines A431 and U2OS (calculated MW of 27.8kDa according to Human NP_006133.1, Mouse NP_061224.2 and Rat XP_003750096.1). Recommended concentration: 0.03-0.1ug/ml. Primary incubation 1 hour at room temperature. Immunohistochemistry: In paraffin embedded Human Skin shows strong nuclear staining in selected cells of the basal layer. Recommended concentration, 2-4ug/ml.	
Expiration Date	12 months from date of receipt.	