


---


## Product Datasheet

### **BIRC7 Antibody (orb1248007)**

<b>Description</b>	BIRC7 Antibody
<b>Species/Host</b>	Goat
<b>Reactivity</b>	Human
<b>Conjugation</b>	Unconjugated
<b>Tested Applications</b>	ELISA, WB
<b>Immunogen</b>	The immunogen for this antibody is: C-RAPVRSVRT
<b>Target</b>	BIRC7
<b>Preservatives</b>	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Form/Appearance</b>	Liquid
<b>Concentration</b>	500 ug/mL
<b>Storage</b>	Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Note</b>	For research use only
<b>Application notes</b>	Peptide ELISA: antibody detection limit dilution 1:4000. Western Blot: Approx 30kDa and 35kDa in lysates of cell line MOLT4, and 30kDa in lysates of Daudi, Jurkat and K562. These bands correspond to the reported isoforms with calculated MW of 32.8kDa according to NP_647478.1 and 30.9kDa according to NP_071444.1. Recommended concentration: 0.3 - 1 ug/ml.
<b>Clonality</b>	Polyclonal
<b>Uniprot ID</b>	<a href="#">Q96CA5</a>
<b>NCBI</b>	<a href="#">NP_071444.1</a> , <a href="#">NP_647478.1</a>
<b>Dilution Range</b>	Peptide ELISA: antibody detection limit dilution 1:4000. Western Blot: Approx 30kDa and 35kDa in lysates of cell line MOLT4, and 30kDa in lysates of Daudi, Jurkat and K562. These bands correspond to the reported isoforms with calculated MW of 32.8kDa according to NP_647478.1 and 30.9kDa according to NP_071444.1. Recommended concentration: 0.3 - 1 ug/ml.
<b>Expiration Date</b>	12 months from date of receipt.

orb1248007 staining (0.3 ug/ml) of MOLT4...



orb1248007 staining (0.5 ug/ml) of Daudi...