


















Product Datasheet

dUTPase/DUT Antibody (orb1248032)

Description	dUTPase/DUT Antibody	
Species/Host	Goat	
Reactivity	Human	
Conjugation	Unconjugated	
Tested Applications	ELISA, IHC, WB	
Immunogen	The immunogen for this antibody is: C-QLRFARLSEHATAPT	
Target	dUTPase / DUT	
Preservatives	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.	
Form/Appearance	Liquid	
Concentration	500 ug/mL	
Storage	Aliquot and store at -20°C. Minimize freezing and thawing.	
Note	For research use only	
Application notes	<p>Peptide ELISA: antibody detection limit dilution 1:16000. Western Blot: Approx 20kDa band observed in lysates of cell lines HEK293 and HeLa (calculated MW of 17.7kDa according to NP_001939.1). Recommended concentration: 0.1-0.3ug/ml. Primary incubation was 1 hour. Immunohistochemistry: Paraffin embedded Human Testis and Tonsil. Recommended concentration: 5ug/ml.</p>	
Clonality	Polyclonal	
MW	Approx 20kDa	
Uniprot ID	P33316	
NCBI	NP_001020420.1 , NP_001939.1 , NP_001020419.1	
Dilution Range	<p>Peptide ELISA: antibody detection limit dilution 1:16000. Western Blot: Approx 20kDa band observed in lysates of cell lines HEK293 and HeLa (calculated MW of 17.7kDa according to NP_001939.1). Recommended concentration: 0.1-0.3ug/ml. Primary incubation was 1 hour. Immunohistochemistry: Paraffin embedded Human Testis and Tonsil. Recommended concentration: 5ug/ml.</p>	
Expiration Date	12 months from date of receipt.	