





---

## Product Datasheet

### NPR3 Antibody (orb1248804)

<b>Description</b>	NPR3 Antibody	
<b>Species/Host</b>	Goat	
<b>Reactivity</b>	Human	orb1248804 (1 ug/ml) staining of Human P...
<b>Conjugation</b>	Unconjugated	
<b>Tested Applications</b>	ELISA, IHC, WB	
<b>Immunogen</b>	The immunogen for this antibody is: C-HRELREDSIRSHFS	
<b>Target</b>	NPR3	
<b>Preservatives</b>	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.	
<b>Form/Appearance</b>	Liquid	orb1248804 (2 ug/ml) staining of paraffi...
<b>Concentration</b>	500 ug/mL	
<b>Storage</b>	Aliquot and store at -20°C. Minimize freezing and thawing.	
<b>Note</b>	For research use only	
<b>Application notes</b>	Peptide ELISA: antibody detection limit dilution 1:16000. Western Blot: Approx 52kDa band observed in Human Placenta lysates (calculated MW of 59.8kDa according to NP_000899.1). Recommended concentration: 1-3ug/ml. Immunohistochemistry: In paraffin embedded Human Kidney shows staining of brush borders in select PCT. Recommended concentration: 2-4ug/ml. Paraffin embedded Human Prostate. Recommended concentration: 10ug/ml.	
<b>Clonality</b>	Polyclonal	orb1248804 Negative Control showing stai...
<b>MW</b>	Approx 52 kDa	
<b>Uniprot ID</b>	<a href="#">P17342</a>	
<b>NCBI</b>	<a href="#">NP_000899.1</a>	
<b>Dilution Range</b>	Peptide ELISA: antibody detection limit dilution 1:16000. Western Blot: Approx 52kDa band observed in Human Placenta lysates (calculated MW of 59.8kDa according to NP_000899.1). Recommended concentration: 1-3ug/ml. Immunohistochemistry: In paraffin embedded Human Kidney shows staining of brush borders in select PCT. Recommended concentration: 2-4ug/ml. Paraffin embedded Human Prostate. Recommended concentration: 10ug/ml.	
<b>Expiration Date</b>	12 months from date of receipt.	