



Product Datasheet

FOXC2 Antibody (orb1248892)

| Description | FOXC2 Antibody |  |
|----------------------------|--|--|
| Species/Host | Goat | |
| Reactivity | Human | |
| Conjugation | Unconjugated | |
| Tested Applications | ELISA, FC, WB | |
| Immunogen | The immunogen for this antibody is: RHAAPYSYDCTKY | |
| Target | FOXC2 | |
| Preservatives | Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing. | |
| Form/Appearance | Liquid | |
| Concentration | 500 ug/mL | |
| Storage | Aliquot and store at -20°C. Minimize freezing and thawing. | |
| Note | For research use only |  orb1248892 (2 ug/ml) staining of paraffi... |
| Application notes | Peptide ELISA: antibody detection limit dilution 1:4000. Western Blot: Approx 35kDa band observed in lysates of cell lines K562 and HeLa, which was successfully blocked by incubation with the immunizing peptide (calculated MW of 38.8kDa according to Human NP_075555.1). Recommended concentration 1-2ug/ml. Primary incubation 1 hour at room temperature. |  orb1248892 (2 ug/ml) staining of paraffi... |
| Clonality | Polyclonal | orb1248892 Flow cytometric analysis of p... |
| Uniprot ID | Q99958 | |
| NCBI | NP_005242.1 | |
| Dilution Range | Peptide ELISA: antibody detection limit dilution 1:4000. Western Blot: Approx 35kDa band observed in lysates of cell lines K562 and HeLa, which was successfully blocked by incubation with the immunizing peptide (calculated MW of 38.8kDa according to Human NP_075555.1). Recommended concentration 1-2ug/ml. Primary incubation 1 hour at room temperature. | |
| Expiration Date | 12 months from date of receipt. | |