





---

## Product Datasheet

### SPI1 Antibody (orb1248960)

<b>Description</b>	SPI1 Antibody	
<b>Species/Host</b>	Goat	
<b>Reactivity</b>	Human	
<b>Conjugation</b>	Unconjugated	
<b>Tested Applications</b>	ELISA, IHC, WB	orb1248960 (0.03 ug/ml) staining of DAUD...
<b>Immunogen</b>	The immunogen for this antibody is: C-DLYQRQTHEYY	
<b>Target</b>	SPI1	
<b>Preservatives</b>	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.	
<b>Form/Appearance</b>	Liquid	
<b>Concentration</b>	500 ug/mL	
<b>Storage</b>	Aliquot and store at -20°C. Minimize freezing and thawing.	
<b>Note</b>	For research use only	orb1248960 (3.75 ug/ml) staining of para...
<b>Application notes</b>	Peptide ELISA: antibody detection limit dilution 1:32000. Western Blot: Approx 40kDa band observed in lysates of DAUDI and K562 (calculated MW of 31.2kDa according to NP_001074016.1 and of 31.1kDa according to NP_003111.2). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Harendza S et al, J Biol Chem. 2000 Jun 30;275(26):19552-9.; PMID: 10867017). Recommended concentration: 0.03-0.1ug/ml. Primary incubation was 1 hour. Immunohistochemistry: Paraffin embedded Human Spleen and Colon (lymphoid aggregate). Recommended concentration: 3.75ug/ml.	
<b>Clonality</b>	Polyclonal	
<b>Uniprot ID</b>	<a href="#">P17947</a>	orb1248960 (3.75 ug/ml) staining of para...
<b>NCBI</b>	<a href="#">NP_001074016.1</a> , <a href="#">NP_003111.2</a>	
<b>Dilution Range</b>	Peptide ELISA: antibody detection limit dilution 1:32000. Western Blot: Approx 40kDa band observed in lysates of DAUDI and K562 (calculated MW of 31.2kDa according to NP_001074016.1 and of 31.1kDa according to NP_003111.2). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Harendza S et al, J Biol Chem. 2000 Jun 30;275(26):19552-9.; PMID: 10867017). Recommended concentration: 0.03-0.1ug/ml. Primary incubation was 1 hour. Immunohistochemistry: Paraffin embedded Human Spleen and Colon (lymphoid aggregate). Recommended concentration: 3.75ug/ml.	