

---

## Product Datasheet

### UXT Antibody (orb1248966)

<b>Description</b>	UXT Antibody	
<b>Species/Host</b>	Goat	
<b>Reactivity</b>	Human	
<b>Conjugation</b>	Unconjugated	
<b>Tested Applications</b>	ELISA, IHC, WB	orb1248966 (0.03 ug/ml) staining of Huma...
<b>Immunogen</b>	The immunogen for this antibody is: C-DHRDKVYEQLAKY	
<b>Target</b>	UXT	
<b>Preservatives</b>	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.	
<b>Form/Appearance</b>	Liquid	
<b>Concentration</b>	500 ug/mL	
<b>Storage</b>	Aliquot and store at -20°C. Minimize freezing and thawing.	orb1248966 (5 ug/ml) staining of paraffi...
<b>Note</b>	For research use only	
<b>Application notes</b>	Peptide ELISA: antibody detection limit dilution 1:64000. Western Blot: Approx 20kDa band observed in Human Kidney and Skeletal Muscle lysates (calculated MW of 19.6kDa according to NP_705582.1). Recommended concentration: 0.01-0.03ug/ml. Immunohistochemistry: In paraffin embedded Thyroid Gland shows staining of cytoplasm in all epithelial cells, but nuclear staining in some of them. Recommended concentration, 5-10ug/ml.	
<b>Clonality</b>	Polyclonal	
<b>Uniprot ID</b>	<a href="#">Q9UBK9</a>	
<b>NCBI</b>	<a href="#">NP_705582.1</a> , <a href="#">NP_004173.1</a>	
<b>Dilution Range</b>	Peptide ELISA: antibody detection limit dilution 1:64000. Western Blot: Approx 20kDa band observed in Human Kidney and Skeletal Muscle lysates (calculated MW of 19.6kDa according to NP_705582.1). Recommended concentration: 0.01-0.03ug/ml. Immunohistochemistry: In paraffin embedded Thyroid Gland shows staining of cytoplasm in all epithelial cells, but nuclear staining in some of them. Recommended concentration, 5-10ug/ml.	
<b>Expiration Date</b>	12 months from date of receipt.	