




---

## Product Datasheet

### NDRG2 Antibody (orb1249248)

<b>Description</b>	NDRG2 Antibody	
<b>Species/Host</b>	Goat	
<b>Reactivity</b>	Human, Mouse, Rat	
<b>Conjugation</b>	Unconjugated	
<b>Tested Applications</b>	ELISA, IHC, WB	
<b>Immunogen</b>	The immunogen for this antibody is: KEAELAARILLDQGQ-C	orb1249248 (0.1 ug/ml) staining of Human...
<b>Target</b>	NDRG2	
<b>Preservatives</b>	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.	
<b>Form/Appearance</b>	Liquid	
<b>Concentration</b>	500 ug/mL	
<b>Storage</b>	Aliquot and store at -20°C. Minimize freezing and thawing.	
<b>Note</b>	For research use only	
<b>Application notes</b>	Peptide ELISA: antibody detection limit dilution 1:16000. Western Blot: Approx 45kDa band observed in Human Brain (Temporal Cortex), Mouse and Rat Brain lysates (calculated MW of 40.8kDa according to NP_963293.1). Recommended concentration: 0.1-0.3ug/ml. Primary incubation was 1 hour. Immunohistochemistry: Paraffin embedded Human Adrenal Gland. Recommended concentration: 3.75ug/ml.	
<b>Clonality</b>	Polyclonal	orb1249248 (0.3 ug/ml) staining of Mouse...
<b>Uniprot ID</b>	<a href="#">Q9UN36</a>	
<b>NCBI</b>	<a href="#">NP_963293.1</a>	
<b>Dilution Range</b>	Peptide ELISA: antibody detection limit dilution 1:16000. Western Blot: Approx 45kDa band observed in Human Brain (Temporal Cortex), Mouse and Rat Brain lysates (calculated MW of 40.8kDa according to NP_963293.1). Recommended concentration: 0.1-0.3ug/ml. Primary incubation was 1 hour. Immunohistochemistry: Paraffin embedded Human Adrenal Gland. Recommended concentration: 3.75ug/ml.	
<b>Expiration Date</b>	12 months from date of receipt.	orb1249248 (3.75 ug/ml) staining of para...