
Product Datasheet

MALT1 Antibody (orb1252063)

Description	MALT1 Antibody
Species/Host	Mouse
Reactivity	Human
Conjugation	Unconjugated
Tested Applications	IF, IHC, WB
Immunogen	A recombinant fragment (aa 701-808) was used as the immunogen for this antibody.
Target	MALT1
Preservatives	PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide
Form/Appearance	Liquid
Concentration	0.2 mg/mL
Storage	Aliquot and Store at 2-8°C. Avoid freeze-thaw cycles.
Note	For research use only
Application notes	IF: 1-2 ug/mlWB: 0.5-1 ug/mlIHC (FFPE): 0.5-1 ug/ml for 30 minutes at RT (1)Prediluted format : incubate for 30 min at RT (2)Titerting of the MALT1 antibody may be required for optimal performance.1. FFPE testing requires sections to be boiled in pH6 10mM citrate buffer for 10-20 minutes, followed by cooling at RT for 20 minutes, prior to staining.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Isotype	IgG1, kappa
Clonality	Monoclonal
Uniprot ID	Q9UDY8
Dilution Range	IF: 1-2 ug/mlWB: 0.5-1 ug/mlIHC (FFPE): 0.5-1 ug/ml for 30 minutes at RT (1)Prediluted format : incubate for 30 min at RT (2)Titerting of the MALT1 antibody may be required for optimal performance.1. FFPE testing requires sections to be boiled in pH6 10mM citrate buffer for 10-20 minutes, followed by cooling at RT for 20 minutes, prior to staining.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Expiration Date	12 months from date of receipt.



Western blot testing of Daudi lysate wit...



IHC staining of tonsil tissue with MALT1...