

---

## Product Datasheet

### F13A1 Antibody (orb1252149)

**Description**

F13A1 Antibody

**Species/Host**

Mouse

**Reactivity**

Human

**Conjugation**

Unconjugated

**Tested Applications**

ELISA, FC, IF, IHC-P, WB

**Immunogen**

A human partial recombinant protein corresponding to amino acids 46-181 was used as the immunogen for the Factor XIIIa antibody.

**Target**

F13A1

**Preservatives**

PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide

**Form/Appearance**

Liquid

**Concentration**

0.2 mg/mL

**Storage**

Aliquot and Store at 2-8°C. Avoid freeze-thaw cycles.

**Note**

For research use only

**Application notes**

ELISA: order the BSA-free formulation for coating  
 Immunofluorescence: 1-2 ug/ml  
 Flow Cytometry: 0.5-1 ug/million cells in 0.1ml  
 Immunohistochemistry (FFPE): 1-2 ug/ml for 30 min at RT  
 Prediluted IHC only format: incubate for 30 min at RT (1)  
 Titering of the Factor XIIIa antibody may be required for optimal performance.  
 1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

**Isotype**

IgG2b, kappa

**Clonality**

Monoclonal


**Uniprot ID**
**P00488**
**Dilution Range**

ELISA: order the BSA-free formulation for coating  
 Immunofluorescence: 1-2 ug/ml  
 Flow Cytometry: 0.5-1 ug/million cells in 0.1ml  
 Immunohistochemistry (FFPE): 1-2 ug/ml for 30 min at RT  
 Prediluted IHC only format: incubate for 30 min at RT (1)  
 Titering of the Factor XIIIa antibody may be required for optimal performance.  
 1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.


**Expiration Date**

12 months from date of receipt.

Human Protein Microarray Specificity Validation




Analysis of HuProt(TM) microarray contain...



IHC testing of FFPE human histiocytoma

W...



SDS-PAGE Analysis of Purified, BSA-Free ...