


Product Datasheet

TFRC Antibody (orb1252162)


Description	TFRC Antibody
Species/Host	Mouse
Reactivity	Human
Conjugation	Unconjugated
Tested Applications	ELISA, FC, IF, IHC-P, WB
Immunogen	Amino acids 94-212 (extracellular domain) of the human protein were used as the immunogen for the Transferrin Receptor / CD71 antibody.
Target	TFRC
Preservatives	PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide
Form/Appearance	Liquid
Concentration	0.2 mg/mL
Storage	Aliquot and Store at 2-8°C. Avoid freez-thaw cycles.
Note	For research use only
Application notes	WB: 0.5-1 ug/ml ELISA: 2-4 ug/ml; order the BSA free formulation for coating Flow Cytometry: 0.5-1 ug/million cells in 0.1ml Immunofluorescence: 1-2 ug/ml IHC (FFPE): 1-2 ug/ml for 30 min at RT Optimal dilution of the CD71 antibody should be determined by the researcher.
Isotype	IgG2b, kappa
Clonality	Monoclonal
Uniprot ID	P02786
Dilution Range	WB: 0.5-1 ug/ml ELISA: 2-4 ug/ml; order the BSA free formulation for coating Flow Cytometry: 0.5-1 ug/million cells in 0.1ml Immunofluorescence: 1-2 ug/ml IHC (FFPE): 1-2 ug/ml for 30 min at RT Optimal dilution of the CD71 antibody should be determined by the researcher.
Expiration Date	12 months from date of receipt.



IHC testing of FFPE human placenta tissu...



Western blot testing of human 1) Jurkat,...



IHC testing of FFPE human bone marrow wi...