
Product Datasheet

MFGE8 Antibody (orb1252491)

Description

MFG8 Antibody

Species/Host

Mouse

Reactivity

Human

Conjugation

Unconjugated

Tested Applications

FC, IF, IHC-P, WB

Immunogen

A delipidated human milk fat globule membrane preparation was used as the immunogen for the Lactadherin antibody.

Target

MFG8

Preservatives

PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide

Form/Appearance

Liquid

Concentration

0.2 mg/mL

Storage

Aliquot and Store at 2-8°C. Avoid freeze-thaw cycles.

Note

For research use only

Application notes

Western blot: 1-2 ug/ml
Flow Cytometry: 0.5-1 ug/million cells in 0.1ml
Immunofluorescence: 1-2 ug/ml
Immunohistochemistry (FFPE): 0.5-1 ug/ml for 30 min at RT
(1) Prediluted format : incubate for 30 min at RT
(2) Optimal dilution of the Lactadherin antibody should be determined by the researcher.
1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Isotype


IgG1, kappa

Clonality


Monoclonal

Uniprot ID
[Q08431](#)
Dilution Range


Western blot: 1-2 ug/ml
Flow Cytometry: 0.5-1 ug/million cells in 0.1ml
Immunofluorescence: 1-2 ug/ml
Immunohistochemistry (FFPE): 0.5-1 ug/ml for 30 min at RT
(1) Prediluted format : incubate for 30 min at RT
(2) Optimal dilution of the Lactadherin antibody should be determined by the researcher.
1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval



Formalin-fixed, paraffin-embedded human ...



FFPE human lung carcinoma tested with La...



FFPE human renal cell carcinoma tested w...