



Product Datasheet

CCL17 Antibody (orb1272297)

Description	CCL17 Antibody	
Species/Host	Rabbit	
Reactivity	Human	
Conjugation	Unconjugated	
Tested Applications	ELISA, NeA, WB	To detect hTARC by Western Blot analysis...
Immunogen	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hTARC (human Thymus and Activation Regulated Chemokine).	
Target	CCL17	
Form/Appearance	Lyophilized	
Concentration	batch dependent	
Storage	Tarc antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.	
Note	For research use only	To detect hTARC by Western Blot analysis...
Clonality	Polyclonal	
Uniprot ID	Q92583	
NCBI	Q92583	
Dilution Range	Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of hTARC (100.00 ng/mL), a concentration of 20.0 - 30 µg/mL of this antibody is required. ELISA:To detect hTARC by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hTARC.Sandwich:To detect hTARC by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Human TARC as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hTARC. Western Blot:To detect hTARC by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hTARC is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.	