


Product Datasheet

Retn Antibody (orb1272330)


Description	Retn Antibody
Species/Host	Goat
Reactivity	Mouse
Conjugation	Unconjugated
Tested Applications	ELISA, WB
Immunogen	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant mResistin (murine Resistin).
Target	Retn
Form/Appearance	Lyophilized
Concentration	batch dependent
Storage	Resistin antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.
Note	For research use only
Clonality	Polyclonal
Uniprot ID	Q99P87
NCBI	Q99P87
Dilution Range	ELISA:To detect mResistin by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant mResistin.Sandwich:To detect mResistin by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Murine Resistinas a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mResistin. Western Blot:To detect mResistin by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant mResistin is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
Expiration Date	12 months from date of receipt.



To detect mResistin by Western Blot anal...



To detect mResistin by sandwich ELISA (u...



To detect mResistin by Western Blot anal...