



Product Datasheet

PPBP Antibody (orb1272355)

Description	PPBP Antibody	
Species/Host	Goat	<p>To detect hNAP-2 by sandwich ELISA (usin...</p>
Reactivity	Human	
Conjugation	Unconjugated	
Tested Applications	ELISA, WB	
Immunogen	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hNAP-2 (human Neutrophil Activating Protein-2).	
Target	PPBP	
Form/Appearance	Lyophilized	
Concentration	batch dependent	
Storage	NAP-2 antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.	
Note	For research use only	<p>To detect hNAP-2 by Western Blot analysi...</p>
Clonality	Polyclonal	
Uniprot ID	P02775	
NCBI	P02775	
Dilution Range	<p>ELISA:To detect hNAP-2 by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hNAP-2. Sandwich:To detect hNAP-2 by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Human NAP-2as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hNAP-2.Western Blot:To detect hNAP-2 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hNAP-2 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.</p>	
Expiration Date	12 months from date of receipt.	<p>To detect hNAP-2 by Western Blot analysi...</p>