




---

## Product Datasheet

### Ccl7 Antibody (Biotin) (orb1272393)

Description	Ccl7 Antibody (Biotin)	Biorbyt MCP-3
<b>Species/Host</b>	Goat	
<b>Reactivity</b>	Mouse	<p>To detect mMCP-3 by sandwich ELISA (usin...</p>
<b>Conjugation</b>	Biotin	
<b>Tested Applications</b>	ELISA, WB	<p>To detect mMCP-3 by Western Blot analysi...</p>
<b>Immunogen</b>	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant mMCP-3 (murine MCP-3).	
<b>Target</b>	Ccl7	<p>To detect mMCP-3 by Western Blot analysi...</p>
<b>Form/Appearance</b>	Lyophilized	
<b>Concentration</b>	batch dependent	
<b>Storage</b>	MCP-3 antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.	
<b>Note</b>	For research use only	
<b>Application notes</b>	<p>ELISA: Sandwich: To detect mMCP-3 by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.25 - 1.0 µg/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our Polyclonal Anti-Murine MCP-3 (XP-5223) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mMCP-3. Western Blot: To detect mMCP-3 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant mMCP-3 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.</p>	
<b>Clonality</b>	Polyclonal	
<b>Uniprot ID</b>	<a href="#">Q03366</a>	
<b>NCBI</b>	<a href="#">Q03366</a>	
<b>Dilution Range</b>	<p>ELISA: Sandwich: To detect mMCP-3 by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.25 - 1.0 µg/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our Polyclonal Anti-Murine MCP-3 (XP-5223) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mMCP-3. Western Blot: To detect mMCP-3 by Western Blot analysis this antibody can be</p>	