


---

## Product Datasheet

### Ifng Antibody (Biotin) (orb1272504)

Description	Ifng Antibody (Biotin)	
<b>Species/Host</b>	Goat	
<b>Reactivity</b>	Rat	
<b>Conjugation</b>	Biotin	
<b>Tested Applications</b>	ELISA, WB	To detect Rat IFN-gamma by sandwich ELIS...
<b>Immunogen</b>	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant Rat IFN-gamma (Rat IFN-gamma).	
<b>Target</b>	Ifng	
<b>Form/Appearance</b>	Lyophilized	
<b>Concentration</b>	batch dependent	
<b>Storage</b>	IFN-gamma antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.	
<b>Note</b>	For research use only	
<b>Application notes</b>	<p>ELISA:Sandwich:To detect Rat IFN-γ by sandwich ELISA (using 100 μL/well antibody solution) a concentration of 0.25 - 1.0 μg/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our Polyclonal Anti-Rat IFN-γ (XP-5157) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat IFN-γ.</p> <p>Western Blot:To detect Rat IFN-γ by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 μg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant Rat IFN-γ is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.</p>	
<b>Clonality</b>	Polyclonal	
<b>Uniprot ID</b>	<a href="#">P01581</a>	
<b>NCBI</b>	<a href="#">P01581</a>	
<b>Dilution Range</b>	<p>ELISA:Sandwich:To detect Rat IFN-γ by sandwich ELISA (using 100 μL/well antibody solution) a concentration of 0.25 - 1.0 μg/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our Polyclonal Anti-Rat IFN-γ (XP-5157) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat IFN-γ.</p> <p>Western Blot:To detect</p>	