
Product Datasheet

CCL14 Antibody (orb1272513)

Description

CCL14 Antibody


Species/Host

Rabbit

Reactivity

Human

Conjugation

Unconjugated

Tested Applications

ELISA, WB

Immunogen

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hHCC-1 human Macrophage-Derived Chemokine).

Target

CCL14

Form/Appearance

Lyophilized

Concentration

batch dependent

Storage

HCC-1 antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.

Note

For research use only

Clonality


Polyclonal

Uniprot ID
[Q16627](#)
NCBI
[Q16627](#)
Dilution Range

ELISA: To detect hHCC-1 by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hHCC-1. Sandwich: To detect hHCC-1 by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Human HCC-1 as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hHCC-1. Western Blot: To detect hHCC-1 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hHCC-1 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Expiration Date

12 months from date of receipt.



This antibody stained formalin-fixed, pa...