


---

## Product Datasheet

### Cxcl1 Antibody (Biotin) (orb1272522)

Description	Cxcl1 Antibody (Biotin)
<b>Species/Host</b>	Rabbit
<b>Reactivity</b>	Rat
<b>Conjugation</b>	Biotin
<b>Tested Applications</b>	ELISA, WB
<b>Immunogen</b>	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant rat GRO/KC (rat GRO/KC).
<b>Target</b>	Cxcl1
<b>Form/Appearance</b>	Lyophilized
<b>Concentration</b>	batch dependent
<b>Storage</b>	GRO/KC antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.
<b>Note</b>	For research use only
<b>Application notes</b>	ELISA: Sandwich: To detect Rat GRO/KC by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.25 - 1.0 µg/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our Polyclonal Anti-Rat GRO/KC (XP-5146) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat GRO/KC. Western Blot: To detect rat GRO/KC by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant rat GRO/KC is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
<b>Clonality</b>	Polyclonal
<b>Uniprot ID</b>	<a href="#">P14095</a>
<b>NCBI</b>	<a href="#">P14095</a>
<b>Dilution Range</b>	ELISA: Sandwich: To detect Rat GRO/KC by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.25 - 1.0 µg/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our Polyclonal Anti-Rat GRO/KC (XP-5146) as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat



To detect Rat GRO/KC by sandwich ELISA (...)