




---

## Product Datasheet

### Cxcl1 Antibody (orb1272523)

Description	Cxcl1 Antibody	Rat GRO/KC
<b>Species/Host</b>	Rabbit	
<b>Reactivity</b>	Rat	To detect Rat GRO/KC by sandwich ELISA (...)
<b>Conjugation</b>	Unconjugated	
<b>Tested Applications</b>	ELISA, NeA, WB	To detect Rat GRO/KC by Western Blot ana...
<b>Immunogen</b>	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant Rat GRO/KC.	
<b>Target</b>	Cxcl1	To detect Rat GRO/KC by Western Blot ana...
<b>Form/Appearance</b>	Lyophilized	
<b>Concentration</b>	batch dependent	
<b>Storage</b>	GRO/KC antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.	
<b>Note</b>	For research use only	
<b>Clonality</b>	Polyclonal	
<b>Uniprot ID</b>	<b>P14095</b>	
<b>NCBI</b>	<b>P14095</b>	
<b>Dilution Range</b>	<p>Neutralization: To yield one-half maximal inhibition [ND50] of the biological activity of Rat GRO/KC (100 ng/mL), a concentration of 5-10 µg/mL of this antibody is required.</p> <p>ELISA: To detect Rat GRO/KC by direct ELISA (using 100 µL/well antibody solution) a concentration of at least 0.5 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant Rat GRO/KC. Sandwich: To detect Rat GRO/KC by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our Biotinylated Anti-Rat GRO/KC as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat GRO/KC. Western Blot: To detect Rat GRO/KC by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant Rat GRO/KC is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.</p>	To detect Rat GRO/KC by Western Blot ana...