



---

## Product Datasheet

### CCL16 Antibody (Biotin) (orb1272651)

Description	CCL16 Antibody (Biotin)	hLEC
<b>Species/Host</b>	Goat	
<b>Reactivity</b>	Human	To detect hLEC by sandwich ELISA (using ...
<b>Conjugation</b>	Biotin	
<b>Tested Applications</b>	WB	
<b>Immunogen</b>	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hLEC. Human LEC specific antibody was purified by affinity chromatography and then biotinylated.	
<b>Target</b>	CCL16	
<b>Form/Appearance</b>	Lyophilized	
<b>Concentration</b>	batch dependent	
<b>Storage</b>	The lyophilized antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-80°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.	 To detect hLEC by Western Blot analysis ...
<b>Note</b>	For research use only	
<b>Application notes</b>	To detect hLEC by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hLEC is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.	
<b>Clonality</b>	Polyclonal	
<b>Uniprot ID</b>	<a href="#">O15467</a>	
<b>NCBI</b>	<a href="#">O15467</a>	
<b>Dilution Range</b>	To detect hLEC by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hLEC is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.	
<b>Expiration Date</b>	12 months from date of receipt.	