




---

## Product Datasheet

### IL31 Antibody (orb1272663)

Description	IL31 Antibody	
<b>Species/Host</b>	Rabbit	To detect hIL-31 by sandwich ELISA (usin...
<b>Reactivity</b>	Human	
<b>Conjugation</b>	Unconjugated	To detect hIL-31 by Western Blot analysi...
<b>Tested Applications</b>	ELISA, WB	
<b>Immunogen</b>	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hIL-31. Human IL-31 specific antibody was purified by affinity chromatography employing immobilized hIL-31 matrix.	To detect hIL-31 by Western Blot analysi...
<b>Target</b>	IL31	
<b>Form/Appearance</b>	Lyophilized	
<b>Concentration</b>	batch dependent	
<b>Storage</b>	The lyophilized antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-80°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.	
<b>Note</b>	For research use only	
<b>Application notes</b>	ELISA:Indirect:To detect IL-31 by indirect ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant IL-31.SandwichTo detect IL-31 by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0 µg/mL of this antibody is required. This antigen affinity purified antibody, in conjunction with our biotinylated Anti-Human IL-31 as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant IL-31. Western Blot:To detect IL-31 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant IL-31 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.	
<b>Clonality</b>	Polyclonal	
<b>Uniprot ID</b>	<a href="#">Q6EBC2</a>	
<b>NCBI</b>	<a href="#">Q6EBC2</a>	
<b>Dilution Range</b>	ELISA:Indirect:To detect IL-31 by indirect ELISA (using 100 µL/well antibody solution) a concentration of 0.5 - 2.0	