

www.biorbyt.com

### **Product Datasheet**

### **RETN Antibody (orb1272664)**

# biorbyt


### **RETN** Antibody

Description <sup>nts.</sup>	RETN Antibody
Species/Host	Mouse
Reactivity	Human
Conjugation	Unconjugated
Tested Applications	ELISA, WB
Immunogen	Resistin antibody is produced in BALB/c mice using highly pure (>98%) recombinant human Resistin as the immunizing antigen.
Target	RETN
Form/Appearance	Lyophilized
Concentration	batch dependent
Storage	Resistin lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. Reconstitute in sterile water to a concentration of 0.1 - 1.0 mg/mL. The antibody is stable for at least six weeks at 2-4°C when reconstituted.
Note	For research use only
Application notes	ELISA:In a sandwich ELISA (assuming 100 $\mu$ L/well), a concentration of 4.0-8.0 $\mu$ g/mL of this antibody will detect at least 200 pg/mL of recombinant human Resistin when used with our biotinylated antigen affinity purified antihuman Resistin as the detection antibody at a concentration of at least 0.5-1.0 $\mu$ g/mL. Western Blot:To detect hResistin by Western Blot analysis this antibody can be used at a concentration of 0.20-0.40 $\mu$ g/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hResistin is 0.20-0.40 $\mu$ g/lane, under non-reducing conditions.
Clonality	Monoclonal
Uniprot ID	Q9HD89
NCBI	Q9HD89
Dilution Range	ELISA:In a sandwich ELISA (assuming 100 $\mu$ L/well), a concentration of 4.0-8.0 $\mu$ g/mL of this antibody will detect at least 200 pg/mL of recombinant human Resistin when used with our biotinylated antigen affinity purified anti-

human Resistin as the detection antibody at a

concentration of at least 0.5-1.0 µg/mL. Western Blot:To detect hResistin by Western Blot analysis this antibody can be used at a concentration of 0.20-0.40  $\mu\text{g/mL}.$  Used in conjunction with compatible secondary reagents the

## www.biorbyt.com



To a sandwich ELISA (assuming 100 ul/wel...



To detect hResistin by Western Blot anal...



To detect hResistin by Western Blot anal...

#### Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: info@biorbyt.com | Phone: +44 (0) 1223 859-353 | Fax: +44 (0)1223 280

68 TW Alexander Drive<br>Research Triangle Park<br>Durham, North Carolina<br>27709. United States

Email: info@biorbyt.com | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558