

Product Datasheet

CXCL14 Antibody (Biotin) (orb1272736)

Description	CXCL14 Antibody (Biotin)	
Species/Host	Rabbit	
Reactivity	Human	
Conjugation	Biotin	
Tested Applications	ELISA, WB	To detect hBRAK by sandwich ELISA (using...
Immunogen	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hBRAK. Human BRAK specific antibody was purified by affinity chromatography and then biotinylated.	
Target	CXCL14	
Form/Appearance	Lyophilized	
Concentration	batch dependent	
Storage	BRAK antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.	
Note	For research use only	
Application notes	ELISA: Sandwich: To detect hBRAK by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.25 - 1.0 µg/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our polyclonal Anti-Human BRAK as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hBRAK. Western Blot: To detect hBRAK by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant hBRAK is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.	
Clonality	Polyclonal	
Uniprot ID	O95715	
NCBI	O95715	
Dilution Range	ELISA: Sandwich: To detect hBRAK by sandwich ELISA (using 100 µL/well antibody solution) a concentration of 0.25 - 1.0 µg/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our polyclonal Anti-Human BRAK as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hBRAK. Western Blot: To detect hBRAK by	