

Product Datasheet

Ccl4 Antibody (Biotin) (orb1272749)

Description	Ccl4 Antibody (Biotin)	
Species/Host	Rabbit	
Reactivity	Mouse	
Conjugation	Biotin	
Tested Applications	ELISA, WB	
Immunogen	Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant mMIP-1 β . Murine MIP-1 β specific antibody was purified by affinity chromatography and then biotinylated.	To detect Murine MIP-1-beta by sandwich ...
Target	Ccl4	
Form/Appearance	Lyophilized	
Concentration	batch dependent	
Storage	MIP-1B antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.	
Note	For research use only	
Application notes	ELISA:Sandwich:To detect mMIP-1b by sandwich ELISA (using 100 μ L/well antibody solution) a concentration of 0.25 - 1.0 μ g/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our polyclonal Anti-Murine MIP-1b as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mMIP-1b. Western Blot:To detect mMIP-1b by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 μ g/mL. Used in conjunction with compatible secondary reagents the detection limit for recombinant mMIP-1b is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.	
Clonality	Polyclonal	
Uniprot ID	P14097	
NCBI	P14097	
Dilution Range	ELISA:Sandwich:To detect mMIP-1b by sandwich ELISA (using 100 μ L/well antibody solution) a concentration of 0.25 - 1.0 μ g/mL of this antibody is required. This biotinylated polyclonal antibody, in conjunction with our polyclonal Anti-Murine MIP-1b as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mMIP-1b. Western Blot:To detect mMIP-1b	