

---

## Product Datasheet

### TNF Antibody (orb1274250)

**Description**

TNF Antibody

**Species/Host**

Rabbit

**Reactivity**

Human

**Conjugation**

Unconjugated

**Tested**

IF, IHC-P

**Applications**
**Immunogen**

Recombinant human protein was used as the immunogen for the TNF-alpha antibody.

**Target**

TNF

**Preservatives**

PBS with 0.1 mg/ml rAlbumin and 0.05% sodium azide

**Form/Appearance**

Liquid

**Concentration**

0.2 mg/mL

**Storage**

Aliquot and Store at 2-8°C. Avoid freeze-thaw cycles.

**Note**

For research use only

**Application notes**

Immunofluorescence: 5-10 ug/ml  
 Immunohistochemistry (FFPE): 5-10 ug/ml for 30 min at RT (1)  
 Optimal dilution of the TNF-alpha antibody should be determined by the researcher.  
 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.

**Isotype**

IgG, k

**Clonality**


Recombinant

**Uniprot ID**
**P01375**
**Dilution Range**


Immunofluorescence: 5-10 ug/ml  
 Immunohistochemistry (FFPE): 5-10 ug/ml for 30 min at RT (1)  
 Optimal dilution of the TNF-alpha antibody should be determined by the researcher.  
 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.

**Expiration Date**

12 months from date of receipt.



Formalin-fixed, paraffin-embedded human ...



Formalin-fixed, paraffin-embedded human ...