
Product Datasheet

CD47 antibody (orb1293382)

Evolution Biocomponents.
Description

Rat monoclonal antibody to CD47

Species/Host	Rat
Reactivity	Mouse
Conjugation	Unconjugated
Tested Applications	FC, IF
Immunogen	Native purified mouse CD47
Target	CD47
Preservatives	Rat IgG2a kappa. Liquid in PBS, pH 7.3, and 0.02% sodium azide.
Form/Appearance	Rat IgG2a kappa. Liquid in PBS, pH 7.3, and 0.02% sodium azide.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Note	For research use only
Application notes	1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5 ml tube. 2. Add appropriate amount of antibody to the bottom of flow tube mixing with the whole blood; incubate for 30 minutes at room temperature. 3. Add 2 ml RBC lysis buffer; incubate for 10 minutes after mixing; dissolve red blood cells. 4. Sample tube is set to 1000 rpm centrifugation for 5 minutes; discard the supernatant. 5. Add 2 ml PBS wash buffer to resuspend the cells; then 1000 rpm centrifugation for 5 minutes; discard the supernatant. 6. Add appropriate amount of fluorescent-labeled anti-rat IgGs and incubate for 20 minutes away from light at room temperature. 7. Add 2 ml PBS wash buffer to resuspend the cells; then 1000 rpm centrifugation for 5 minutes; discard the supernatant. 8. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).
Clonality	Monoclonal
Source	Rat
Uniprot ID	Q61735