

Product Datasheet

RecombinantIL-7, His, Rat (orb1494693)

Description	Interleukin-7 (IL-7), also known as lymphopoietin 1 and pre-B cell factor, is a hematopoietic growth factor belonging to the IL-7/IL-9 family. It is produced by keratinocytes, dendritic cells, hepatocytes, neurons and epithelial cells. IL-7 binds and signals through IL-7 receptor, a heterodimer consisting of IL-7 receptor alpha and common gamma chain receptor. IL-7 plays a role in regulating early B cell and T cell development. It is also important for optimal dendritic cell-T cell interaction.
Endotoxins	< 0.2 EU/ μ g, determined by LAL method.
Preservatives	Lyophilized after extensive dialysis against PBS.
Form/Appearance	Lyophilized after extensive dialysis against PBS.
Storage	Lyophilized recombinant rat Interleukin-7 (IL-7), His remains stable up to 6 months at -80°C from date of receipt. Upon reconstitution, rat Interleukin-7 (IL-7), His should be stable up to 1 week at 4°C or up to 2 months at -20°C.
Note	For research use only
Application notes	Reconstituted in ddH2O or PBS at 100 μ g/ml.
Protein Sequence	DCHIKDKDGKAFGSVLMISINQLDKMTGTDSDCPNNEPNFFKKHLCDDTKEAAFLNRAARK LRQFLKMNISEEFNDHLLR VSDGTQTLVNCTSKEEKTIKEQKKNDPCFLKRLLREIKTCWNKILKGSIHHHHHH
Purity	> 95% as analyzed by SDS-PAGE.
Source	СНО
MW	20-28 kDa, observed by reducing SDS-PAGE.
Expiration Date	6 months from date of receipt.

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +44 (0) 1223 859-353 | Fax: +1 (415) 651-8558

Biorbyt LLC.

68 TW Alexander Drive, Durham, NC, 27713, United States Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>