

Product Datasheet

RecombinantIL-4, His, Rat (orb1494706)

Description	Interleukin-4 (IL-4), also known as BSF-1 and Lymphocyte stimulatory factor 1, is a secreted cytokine belonging to the IL-4/IL-13 family. It is produced by mast cells, T cells and bone marrow stromal cells. IL-4 signals through two receptor complexes: theIL4R α /common γ chain and the IL4R α -IL-13 R α 1 receptor complex. IL-4 regulates T cell and B cell proliferation, survival and gene expression. IL-4 also induces IgG and IgE class switching, and the upregulation of MHC-II production.
Endotoxins	< 0.2 EU/ μ g, determined by LAL method.
Preservatives	Lyophilized after extensive dialysis against PBS.
Form/Appearance	Lyophilized after extensive dialysis against PBS.
Storage	Lyophilized recombinant rat Interleukin-4 (IL-4), Hisremains stable up to 6 months at -80°C from date of receipt. Upon reconstitution, rat Interleukin-4 (IL- 4), Hisshould be stable up to 1 week at 4°C or up to 2 months at -20°C.
Note	For research use only
Application notes	Reconstituted in ddH2O or PBS at 100 μ g/ml.
Protein Sequence	HGCNDSPLREIINTLNQVTEKGTPCTEMFVPDVLTATRNTTENELICRASRVLRKFYFPRDVP PCLKNKSGVLGELRKLC RGVSGLNSLRSCTVNESTLTTLKDFLESLKSILRGKYLQSCTSMSHHHHHH
Purity	> 95% as analyzed by SDS-PAGE.
Source	СНО
MW	18-22 kDa, observed by reducing SDS-PAGE.
Expiration Date	6 months from date of receipt.

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: +44 (0) 1223 859-353 | Fax: +1 (415) 651-8558

Biorbyt LLC.

68 TW Alexander Drive, Durham, NC, 27713, United States Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>