

Product Datasheet

RecombinantIL-10,Rat(CHO-expressed) (orb1494721)

Description Interleukin-10 (IL-10), initially known as Cytokine Synthesis Inhibitory Factor

(CSIF), belongs to the IL-10 family and shares more than 80% sequence homology with the Epstein-Barr Virus protein BCRFI. It is produced by many immune cells, such as T-cells, macrophages, mast cells and dendritic cells. It is usually secreted as a homodimer and, upon binding to its receptor, inhibits the synthesis of a number of cytokines, including IFN-gamma, IL-2, IL-3, TNF and GM-CSF produced by activated macrophages and Th2 cells. It also displays the ability to suppress Antigen-Presenting Cell (APC) function. The net effect of Interleukin-10 appears to be inhibitory; however, stimulatory effects, such as stimulation of B cell maturation and antibody production, are also reported.

Endotoxins $< 0.2 \text{ EU/}\mu\text{g}$, determined by LAL method.

Preservatives Lyophilized after extensive dialysis against PBS.

Form/Appearance Lyophilized after extensive dialysis against PBS.

Storage Lyophilized recombinant rat Interleukin-10(IL-10) remains stable up to 6 months

at -80°C from date of receipt. Upon reconstitution, Rat Interleukin-10 should be

stable up to 1 week at 4°C or up to 2 months at -20°C.

Note For research use only

Application notes Reconstituted in ddH2O or PBS at 100 μg/ml.

Protein Sequence SKGHSIRGDNNCTHFPVSQTHMLRELRAAFSQVKTFFQKKDQLDNILLTDSLLQDFKGYLG

CQALSEMIKFYLVEVMPQA

ENHGPEIKEHLNSLGEKLKTLWIQLRRCHRFLPCENKSKAVEQVKNDFNKLQDKGVYKAMN

EFDIFINCIEAYVTLKMKN

Purity > 95% as analyzed by SDS-PAGE and HPLC.

Source CHO

MW 8-22 kDa, observed by reducing SDS-PAGE.

Biorbyt Ltd.

Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558





Expiration Date

6 months from date of receipt.

68 TW Alexander Drive, Durham, NC, 27713, United States

Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u>
Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>