

---

## Product Datasheet

### **Murine Stromal-Cell Derived Factor-1 alpha/CXCL12 (rMuSDF-1a/CXCL12) (orb1495005)**

**Description**

SDF-1α and SDF-1β, members

that lack the ELR domain, were initially ...

**Endotoxins**

Less than 1EU/μg of rMuSDF-1a/CXCL12 as determined by LAL method.

**Preservatives**

Lyophilized from a 0.2mm filtered concentrated solution in PBS, pH 7.4.

**Form/Appearance**

Sterile Filtered White lyophilized (freeze-dried) powder.

**Storage**

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

**Note**

For research use only

**Application notes**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at -20°C. Further dilutions should be made in appropriate buffered solutions.

**Protein Sequence**

KPVLSYRCP CRFFESHIAR  
ANVKHLKILN TPNCALQIVA  
RLKNNNRQVC IDPKLKWIQE  
YLEKALNK

**Source**

Escherichia coli.

**MW**

Approximately 7.9 kDa, a single non-glycosylated polypeptide chain containing 68 amino