

## Product Datasheet

# Recombinant Bovine Fibroblast Growth Factor-basic (rBoFGF-2) (orb1495071)

<b>Description</b>	bFGF is a single-chain polypeptide growth factor that plays a significant role in the process of wound healing and is a potent inducer of angiogenesis. Several different forms of the human protein exist ranging from 18-24 kDa in size due to the use of alternative start sites within the FGF-2 gene. It has a 55 percent amino acid residue identity to FGF-1 and has potent heparin-binding activity. The growth factor is an extremely potent inducer of DNA synthesis in a variety of cell types from mesoderm and neuroectoderm lineages. It was originally named basic fibroblast growth factor based upon its chemical properties and to distinguish it from acidic fibroblast growth factor.
<b>Endotoxins</b>	Less than 1EU/mg of rBoFGF-2 as determined by LAL method.
<b>Preservatives</b>	Lyophilized from a 0.2mm filtered concentrated solution in PBS, pH 7.4, containing 4% mannitol.
<b>Form/Appearance</b>	Lyophilized from a 0.2mm filtered concentrated solution in PBS, pH 7.4, containing 4% mannitol.
<b>Storage</b>	This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.
<b>Note</b>	For research use only
<b>Application notes</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at -20°C. Further dilutions should be made in appropriate buffered solutions.

### Biorbyt Ltd.

7 Signet Court, Swann's Road,  
Cambridge, CB5 8LA, United Kingdom  
Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

### Biorbyt LLC.

68 TW Alexander Drive,  
Durham, NC, 27713, United States  
Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

<b>Protein Sequence</b>	MTMITNSSSV PGDPLESMAS GSITTLPALP EDGGSGAFPP GHFKDPKRLY CKNGGFFLRI HPDGRVDGVR EKSDPHIKLQ LQAEERGVVS IKGVCANRYL AMKEDGRLLA SKCVTDECFF FERLESNNYN TYRSRKYSSW YVALKRTGQY KLGPKTGPGQ KAILFLPMSA KS
<b>Purity</b>	> 95% by SDS-PAGE and HPLC analyses.
<b>Source</b>	Escherichia coli.
<b>MW</b>	Approximately 18.5 kDa, a single non-glycosylated polypeptide chain containing 172 amino acids
<b>Expiration Date</b>	6 months from date of receipt.

**Biorbyt Ltd.**

7 Signet Court, Swann's Road,  
Cambridge, CB5 8LA, United Kingdom  
Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

**Biorbyt LLC.**

68 TW Alexander Drive,  
Durham, NC, 27713, United States  
Email: [info@biorbyt.com](mailto:info@biorbyt.com), [support@biorbyt.com](mailto:support@biorbyt.com)  
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)