

Product Datasheet

Mouse IgG1 Isotype control (APC) (orb154438)

Description

Mouse IgG1 Isotype control conjugated to APC which



Flow cytometry: Example of nonspecific m...

Conjugation

APC

Tested Applications

FC

Preservatives

Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide

Concentration

0.1 mg/ml

Storage

Store at 2-8°C. Protect from prolonged exposure to light. Do not freeze.

Note

For research use only

Application notes

The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions

Isotype

Mouse IgG1 kappa

Clonality

Monoclonal

Purity

Purified antibody is conjugated with activated allophycocyanin (APC) under optimum conditions and unconjugated antibody and free fluorochrome are removed by size-exclusion chromatography.

Dilution Range

Negative control: The reagent is intended as an isotype control to establish the amount of non-specific antibody binding. For your particular experiment, use the same concentration of this control antibody as the recommended working concentration of the antigen-specific antibody. Also, when working with prediluted antibodies, dilute the isotype control to the same concentration as is the concentration of the antigen-specific antibody in the prediluted antibody solution you are using. If under particular experimental conditions the background signal of the isotype control is too high (usually when working concentrations of used antibodies are above 10 µg/ml of incubation mixture), change the conditions of your experiment to reduce the background.

Expiration Date

12 months from date of receipt.