

---


## Product Datasheet

### CCL15 Antibody (orb1671450)

<b>Description</b>	CCL15 Antibody
<b>Species/Host</b>	Goat
<b>Reactivity</b>	Human
<b>Conjugation</b>	Unconjugated
<b>Tested Applications</b>	ELISA, WB
<b>Immunogen</b>	Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hMIP-5. Anti-Human MIP-5 specific antibody was purified by affinity chromatography employing immobilized hMIP-5 matrix.
<b>Target</b>	CCL15
<b>Form/Appearance</b>	Lyophilized
<b>Concentration</b>	batch dependent
<b>Storage</b>	Antibody is stable for at least 2 years from date of receipt at -20°C. The reconstituted antibody is stable for at least two weeks at 2-8°C. Frozen aliquots are stable for at least 6 months when stored at -20°C. Avoid repeated freeze-thaw cycles.
<b>Note</b>	For research use only
<b>Application notes</b>	Sandwich ELISA: To detect hMIP-5 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with Biorbyt's Biotinylated Anti-Human MIP-5 as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMIP-5. Western Blot To detect hMIP-5 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 ug/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMIP-5 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
<b>Clonality</b>	Polyclonal
<b>Uniprot ID</b>	<a href="#">Q16663</a>
<b>NCBI</b>	<a href="#">Q16663</a>
<b>Expiration Date</b>	12 months from date of receipt.



To detect hMIP-5 by sandwich ELISA (usin...



To detect hMIP-5 by Western Blot analysi...