
Product Datasheet

NR1H2 antibody (orb18918)

Description

Goat polyclonal antibody to NR1H2


Species/Host

Goat

Reactivity

Canine, Human, Mouse, Rat

Conjugation

Unconjugated

Tested

ELISA, FC, IF

Applications

Immunohistochemical staining of paraffin...

Target

LXR beta / NR1H2

Preservatives

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

Storage

Aliquot and store at -20°C. Minimize freezing and thawing.

Note

For research use only

Application notes

ELISA: Peptide ELISA: antibody detection limit dilution 1:64000. IHC: In paraffin embedded Human Spleen shows nucleolar staining. Recommended concentration, 1-2ug/ml. WB: Experiments gave a band at approx 80kDa and a weak band at appr. 60-kDa in human bone marrow lysate after 1 µg/ml antibody staining. These bands correspond to earlier findings with different antibodies (PMID: 12393874-60kDa and other unpublished antibodies-80kDa). This protein has a calculated MW of 51.0kDa according to NP_009052. Recommended concentration: 1-3 µg/ml.

Immunohistochemical staining of Human Ut...



Western blot analysis of Human Bone Marr...

Protein Sequence

SSPTTSSLDTPLPGC

Clonality

Polyclonal

MW

51

Entrez
7376
Dilution Range

ELISA: 1:128000, IHC-P: 10ug/ml

Expiration Date

12 months from date of receipt.