



Product Datasheet

Munc13-4 antibody (orb19314)


Description	Goat polyclonal to Munc13-4
Species/Host	Goat
Reactivity	Bovine, Canine, Human, Mouse, Porcine, Rat
Conjugation	Unconjugated
Tested Applications	ELISA, IHC, WB
Target	Munc13-4 / UNC13D (Internal)
Preservatives	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Storage	Aliquot and store at -20°C. Minimize freezing and thawing.
Note	For research use only
Application notes	ELISA: Peptide ELISA: antibody detection limit dilution 1:128000.IHC: In paraffin embedded Human Thymus shows cytoplasm staining in select areas. Recommended concentration, 2.5ug/ml.Full caption (WB): EB067336 (2ug/ml) staining of Human T-lymphocyte lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. DeteWB: Approx. 105kDa band observed in Human T-lymphocyte and HeLa lysates (calculated MW of 123kDa according to NP_954712.1). Recommended concentration: 1-3 µg/ml.
Protein Sequence	ETQKHKKDLHPLFD
Clonality	Polyclonal
MW	123
Entrez	201294
Dilution Range	ELISA: 1:128000, WB: 1-3 µg/ml, IHC-P: 1-2 µg/ml
Expiration Date	12 months from date of receipt.



Western blot analysis of Human T-lymphoc...



Immunohistochemical staining of paraffin...



Immunohistochemical staining of Mouse Lu...