

Product Datasheet

CD130 Antibody (C-term) (orb1938446)

Description	Affinity Purified Rabbit Polyclonal Antibody (Pab)
Species/Host	Rabbit
Reactivity	Human
Conjugation	Unconjugated
Tested Applications	IHC-P, WB
Target	This CD130 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 871-899 amino acids from the C-terminal region of human CD130.
Form/Appearance	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles
Note	For research use only
Isotype	Rabbit IgG
Clonality	Polyclonal
Clone Number	RB17993
MW	103537 Da
Uniprot ID	P40189
NCBI	NP_001177910.1, NP_002175.2, NP_786943.1
Dilution Range	WB: 1:2000, WB: 1:2000, WB: 1:2000, WB: 1:1000, WB: 1:2000, WB: 1:2000, IHC-P: 1:25


Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)


Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)


Expiration Date 12 months from date of receipt.



All lanes: Anti-CD130 Antibody (C-term) at 1:2000 dilution. Lane 1: A549 whole cell lysate. Lane 2: HeLa whole cell lysate. Lane 3: HepG2 whole cell lysate. Lane 4: HT-1080 whole cell lysate. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 104kDa. Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes: Anti-CD130 Antibody (C-term) at 1:2000 dilution. Lane 1: A549 whole cell lysate. Lane 2: HepG2 whole cell lysate. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 104 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.




All lanes: Anti-CD130 Antibody (C-term) at 1:2000 dilution. Lane 1: HeLa whole cell lysate. Lane 2: MDA-MB-231 whole cell lysate. Lane 3: HepG2 whole cell lysate. Lane 4: SW480 whole cell lysate. Lane 5: A549 whole cell lysate. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 104 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

Biorbyt Ltd.


7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Biorbyt LLC.


68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)



All lanes: Anti-CD130 Antibody (C-term) at 1:2000 dilution. Lane 1: HepG2 whole cell lysate. Lane 2: MDA-MB-231 whole cell lysate. Lane 3: SW480 whole cell lysate. Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 104 kDa. Blocking/Dilution buffer: 5% NFDm/TBST.



Anti-CD130 Antibody (C-term) at 1:2000 dilution + SW480 whole cell lysates. Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 104 kDa. Blocking/Dilution buffer: 5% NFDm/TBST.




CD130 Antibody (C-term) western blot analysis in MDA-MB231 cell line lysates (35 μ g/lane). This demonstrates the CD130 antibody detected the CD130 protein (arrow).

Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)



Staining CD130 in Human skeletal muscle tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary Antibody.

Biorbyt Ltd.

7 Signet Court, Swann's Road,
Cambridge, CB5 8LA, United Kingdom
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+44 \(0\) 1223 859-353](tel:+44(0)1223859353) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)

Biorbyt LLC.

68 TW Alexander Drive,
Durham, NC, 27713, United States
Email: info@biorbyt.com, support@biorbyt.com
Phone: [+1 \(415\) 906-5211](tel:+1(415)9065211) | Fax: [+1 \(415\) 651-8558](tel:+1(415)6518558)