Product Datasheet

Endostatin Protein, Mouse, Recombinant (His) (orb1961368)

Description	Endostatin, an endogenous non-glycosylated inhibitor of endothelial cell proliferation and angiogenesis. It is produced and/or trimmed by metalloproteinases such as MMP-2 and MMP-9, and cathepsins S, B and L. The N-terminal ~27 aa of Endostatin appear to contain the majority of its activity. This region contains zinc binding sites that are thought to be critical for its anti-endothelial and anti-tumor effects, as well as multiple cleavage sites that, when used, can modify its activity. Mouse Endostatin shares 96% aa sequence identity with rat and 85-87% with human, bovine and equine Endostatin. It is predominantly expressed in liver, kidney, lung, skeletal muscle and testis. Endostatin inhibits endothelial cell growth by inducing cell cycle arrest in G1 phase and initiating apoptosis. It is also thought to down-regulate angiogenesis by blocking VEGF-induced endothelial cell migration. Endostatin may also be involved with down-regulation of angiogenesis after establishment of placental circulation in the pregnant uterus.
Storage	-20°C
Tag	C-6xHis
Note	For research use only
Application notes	Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.
Protein Sequence	His1591-Lys1774
Purity	98.00%
MW	18 KDa (reducing condition)
Uniprot ID	P39061
Expiration Date	6 months from date of receipt.