

Product Datasheet

Sheep anti Guinea Pig IgG (Fc specific), conjugated with Horseradish peroxidase (orb21568)

Description

Horseradish peroxidase-conjugated IgG fraction of polyclonal sheep antiserum to guinea pig IgG, Fc specific. In enzyme-immunocytochemical and immunohistochemical staining for the detection of IgG at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to demon- strate circulating IgG antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen using a reference antibody of guinea pig origin known to be of the IgG isotype in the middle layer of the indirect test procedure; in non-isotopic assay methodology (e.g. ELISA) to measure IgG in guinea pig serum or other body fluids. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions for histochemical and cytochemical use are usually between 1:100 and 1:250; in ELISA and comparable non-precipitating antibody-binding assays between 1:1,000 and 1:5,000.

Species/Host Sheep

Conjugation HRP

Preservatives Horseradish peroxidase-coupled purified hyperimmune sheep IgG lyophilized

from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative

added, as it may interfere with the antibody activity.

Form/Appearance Horseradish peroxidase-coupled purified hyperimmune sheep IgG lyophilized

from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative

added, as it may interfere with the antibody activity.

Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558



Biorbyt.com

Storage

The lyophilized conjugate is shipped at ambient temperature and may be stored at $+4^{\circ}$ C; prolonged storage at or below -20° C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at $+4^{\circ}$ C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate

Note For research use only

Clonality Polyclonal

Source Purified normal IgG isolated from pooled guinea pig serum. Freund's complete

adjuvant is used in the first step of the immunization procedure

Hazard Information This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO,

not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Biorbyt accepts no

liability for any inaccuracies or omissions in this information.

Expiration Date 12 months from date of receipt.