

## **Product Datasheet**

## Rabbit anti Monkey IgG IgA IgM (heavy and light chains), conjugated with Biotin (orb21722)

**Description** 

Biotin-conjugated IgG fraction of polyclonal goat antiserum to monkey IgG, IgA and IgM, heavy and light chains. In immunocytochemical and immunohistochemical staining of immunoglobulins at the cellular and subcellular level of appropriately treated cell and tissue substrates; to demonstrate circulating antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen using a reference antibody of monkey origin in the middle layer of the indirect test procedure; in non-isotopic assay methodology (e.g. ELISA) to measure immunoglobulins in monkey serum or other body fluids. As a second step an avidin or streptavidin conjugate of the user's choice has to be used. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions for histochemical and cytochemical use are usually between 1:100 and 1:500; in ELISA and comparable non-precipitating antibody-binding assays between 1:2,000 and 1:20,000.

Species/Host Rabbit

**Conjugation** Biotin

**Preservatives**Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may

interfere with the antibody activity.

**Form/Appearance** Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in

phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may

interfere with the antibody activity.

Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>



## **Biorbyt.com**

**Storage** 

The lyophilized conjugate is shipped at ambient temperature and may be stored at  $+4^{\circ}$ C; prolonged storage at or below  $-20^{\circ}$ C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at  $+4^{\circ}$ C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.

**Note** For research use only

**Clonality** Polyclonal

**Source** Highly purified IgG, IgA and IgM isolated from Rhesus monkey serum. Freund's

complete adjuvant is used in the first step of the immunization procedure.

**Hazard Information** This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO,

not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Biorbyt accepts no

liability for any inaccuracies or omissions in this information.

**Expiration Date** 12 months from date of receipt.