

Product Datasheet

Goat anti Human IgG IgA IgM (heavy and light chains), conjugated with Horseradish peroxidase (orb21831)

Description	Horseradish peroxidase-conjugated IgG fraction of polyclonal goat antiserum to human IgG, IgA and IgM, heavy and light chains. In enzyme- immunocytochemical and immunohistochemical staining for the detection of cytoplasmic Ig at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates, and to demonstrate circulating antibodies in serodiagnostic microbiology and autoimmune diseases. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions for histochemical and cytochemical use are usually between 1:100 and 1:250; in ELISA and comparable non- precipitating antibody-binding assays between 1:1,500 and 1:15,000.
Species/Host	Goat
Reactivity	Human
Conjugation	HRP
Preservatives	Peroxidase-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.
Form/Appearance	Peroxidase-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+44 (0) 1223 859-353</u> | Fax: <u>+1 (415) 651-8558</u>

Biorbyt LLC.

68 TW Alexander Drive, Durham, NC, 27713, United States Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>



Biorbyt.com

Storage	The lyophilized conjugate is shipped at ambient temperature and may be stored at $+4^{\circ}$ C; prolonged storage at or below -20° C. Spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20° C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at $+4^{\circ}$ C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.
Note	For research use only
Clonality	Polyclonal
Source	Purified polyclonal human IgG, and IgA and IgM containing factions isolated from human serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Hazard Information	This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Biorbyt accepts no liability for any inaccuracies or omissions in this information.
Expiration Date	12 months from date of receipt.

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+44 (0) 1223 859-353</u> | Fax: <u>+1 (415) 651-8558</u>

Biorbyt LLC.

68 TW Alexander Drive, Durham, NC, 27713, United States Email: <u>info@biorbyt.com</u>, <u>support@biorbyt.com</u> Phone: <u>+1 (415) 906-5211</u> | Fax: <u>+1 (415) 651-8558</u>