

Product Datasheet

Goat IgG Rat IgG1 IgG2A IgG2B IgG2C IgA IgM (TRITC) antibody (orb22096)



www.biorbyt.com

Descriptionnts. Tetramethylrhodamine

fraction of polyclonal goat antiserum to rat IgG,...

Species/Host Goat

TRITC Conjugation

Preservatives TRITC-coupled purified

> hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.

TRITC-coupled purified Form/Appearance

> hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.

Storage Storage: The lyophilized conjugate

> is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient

temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C. not refrozen, and preferably used

the same day. If a slight

precipitation occurs upon storage,

this should be removed by

centrifugation. It will not affect the

performance of the immunoconjugate

Note For research use only

Clonality Polyclonal

Purified polyclonal rat IgG, and Source

> homogenous IgA and IgM isolated from rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.

Biorbyt Ltd.

7 Signet Court, Swann's Road, Cambridge, CB5 8LA, United Kingdom

Email: info@biorbyt.com | Phone: +44 (0) 1223 859-353 | Fax: +44 (0) 1223 280

Biorbyt LLC.

68 TW Alexander Drive
br>Research Triangle Park
br>Durham, North Carolina < br > 27709. United States

Email: info@biorbyt.com | Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558