



Product Datasheet

VDAC/Porin/VDAC1 Antibody (orb259632)

Description	VDAC/Porin/VDAC1 Antibody	
Species/Host	Rabbit	
Reactivity	Human, Mouse, Rat	
Conjugation	Unconjugated	
Tested Applications	IHC, WB	WB analysis of VDAC using anti-VDAC anti...
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human VDAC/Porin (154-181aa QMNFETAKSRVTQSNFAVGKYKTDEFQLH), different from the related mouse and rat sequences by one amino acid.	
Form/Appearance	Lyophilized	
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.	
Storage	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.	
Note	For research use only	IHC analysis of VDAC using anti-VDAC ant...
Application notes	WB: The detection limit for VDAC/Porin is approximately 0.1ng/lane under reducing conditions. Tested Species: In-house tested species with positive results. By Heat: Boiling the paraffin sections in 10mM citrate buffer, pH6.0, for 20mins is required for the staining of formalin/paraffin sections. Other applications have not been tested. Optimal dilutions should be determined by end users. . Add 0.2ml of distilled water will yield a concentration of 500ug/ml.	
Isotype	Rabbit IgG	
Clonality	Polyclonal	
MW	30773 MW	
Uniprot ID	P21796	
Expiration Date	12 months from date of receipt.	IHC analysis of VDAC using anti-VDAC ant...