

Product Datasheet

Angiotensin Converting Enzyme Antibody / ACE (orb2635152)

Description This gene encodes an enzyme involved in catalyzing the conversion of

angiotensin I into a physiologically active peptide angiotensin II. Angiotensin II is a potent vasopressor and aldosterone-stimulating peptide that controls blood pressure and fluid-electrolyte balance. This enzyme plays a key role in the reninangiotensin system. Many studies have associated the presence or absence of a 287 bp Alu repeat element in this gene with the levels of circulating enzyme or cardiovascular pathophysiologies. Two most abundant alternatively spliced variants of this gene encode two isozymes - the somatic form and the testicular form that are equally active. Multiple additional alternatively spliced variants have been identified but their full length nature has not been determined.

Species/Host Mouse

Reactivity Human

Conjugation Unconjugated

Tested Applications ELISA, IHC-P

Immunogen Full-length recombinant human protein was used as the immunogen for the

Angiotensin Converting Enzyme antibody.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -

20°C in small aliquots to prevent freeze-thaw cycles.

Note For research use only

Application notes Optimal dilution of the Angiotensin Converting Enzyme antibody should be

determined by the researcher.

Formula 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide

Isotype Mouse IgG1, kappa

Clonality Monoclonal

Biorbyt Ltd.





Clone Number ACE/3764

Antibody Type Primary Antibody


Uniprot ID P12821

Hazard Information This Angiotensin Converting Enzyme antibody is available for research use only.


Dilution Range ELISA (order BSA-free format for coating),Immunohistochemistry (FFPE): 1-

2ug/ml

Expiration Date 12 months from date of receipt.




IHC staining of FFPE human colon carcinoma tissue with Angiotensin Converting Enzyme antibody (clone ACE/3764) at 2 ug/ml in PBS for 30 min RT. HIER: boil tissue sections in pH9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human kidney tissue with Angiotensin Converting Enzyme antibody (clone ACE/3764) at 2 ug/ml in PBS for 30 min RT. HIER: boil tissue sections in pH9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.








SDS-PAGE analysis of purified, BSA-free Angiotensin Converting Enzyme antibody (clone ACE/3764) as confirmation of integrity and purity.

Human Protein Microarray Specificity Validation



Analysis of HuProt (TM) microarray containing more than 19000 full-length human proteins using Angiotensin Converting Enzyme antibody (clone ACE/3764). These results demonstrate the foremost specificity of the ACE/3764 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt (TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt (TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.

Phone: +1 (415) 906-5211 | Fax: +1 (415) 651-8558